arXiv:1005.4170v2 [math.OA] 26 May 2010

On MF Property of Reduced Amalgamated Free Products of UHF Algebras

Qihui Li Junhao Shen*

May 27, 2010

Abstract

In this paper, we concentrate on the MF property of reduced free products of unital C*-algebras with amalgamation over finite dimensional C*-algebras. More specifically, we give a necessary and sufficient condition for a reduced free product of two UHF algebras amalgamated over a finite-dimensional C*-algebra with respect to trace preserving conditional expectations to be MF.

1 Introduction

The concept of MF algebras was first introduced by Blackadar and Kirchberg in [4]. If a separable C*- algebra $\mathcal{A}$ can be embedded into $\prod_k \mathcal{M}_{n_k}(\mathbb{C}) / \sum_k \mathcal{M}_{n_k}(\mathbb{C})$ for a sequence of positive integers $\{n_k\}_{k=1}^{\infty}$, then $\mathcal{A}$ is an MF algebra. Many properties of MF algebras were discussed in [4]. For example, it was shown there that an inductive limit of MF algebras is an MF algebra and every subalgebra of an MF algebra is an MF algebra. This class of C*-algebras is of interest for many reasons. For example, it plays an important role in the classification of C*-algebras and it is connected to the question whether the Ext semigroup of a unital separable C*-algebra is a group. This notion is also closely connected to Voiculescu's topological free entropy dimension for a family of self-adjoint elements $x_1, \cdots, x_n$ in a unital C*-algebra $\mathcal{A}$ [22].

In [23], Voiculescu introduced a notion of reduced amalgamated free product of C*-algebras in the framework of his free probability theory. The free product construction is everywhere in the C*-literature these days, and plays an important role in the recent study of C*-algebras [17], [11], [12]. In [13], Haagerup and Thorbjϕrnsen solved a long standing open problem by showing that Ext($C_r^*(F_2)$) is not a group, where $C_r^*(F_2)$ is the reduced C*-algebra of the free group F_2. One remarkable ingredient in their proof is their work on proving that $C_r^*(F_2)$ is an MF algebra. From a quick fact that $(C_r^*(F_2), \tau_{F_2}) =$

---

*The research of second author is partially sponsored by an NSF grant.

$(C_r^*(\mathbb{Z}), \tau_{\mathbb{Z}}) *_{red} (C_r^*(\mathbb{Z}), \tau_{\mathbb{Z}})$ where τ_{F_2}, $\tau_{\mathbb{Z}}$ are canonical tracial states on $C_r^*(F_2)$ and $C_r^*(\mathbb{Z})$ respectively, one should naturally ask when a reduced free product of two unital MF algebras is MF again. Recently, it was shown in [16] that a reduced free product of finite-dimensional C*-algebras with respect to faithful tracial states is MF. More generally, the authors proved that a reduced free product of unital separable AH algebras with respect to faithful tracial states is an MF algebra.

In this paper, we discuss the MF property of reduced amalgamated free products of unital C*-algebras. We investigate the reduced amalgamated free products of finite-dimensional C*-algebras with respect to trace preserving conditional expectations and obtain the following results.

Theorem 3.2 Let $\mathcal{D} \subseteq \mathcal{M}_k(\mathbb{C})$ be a unital inclusion of C*-algebras where $\mathcal{D}$ is a finite-dimensional abelian C^*-algebra and $E : \mathcal{M}_k(\mathbb{C}) \to \mathcal{D}$ be the trace-preserving conditional expectation from $\mathcal{M}_k(\mathbb{C})$ onto $\mathcal{D}$. Let $(\mathcal{M}_k(\mathbb{C}), E) \underset{\mathcal{D}}{*} (\mathcal{M}_k(\mathbb{C}), E)$ be the reduced amalgamated free product of $\mathcal{M}_k(\mathbb{C})$ and $\mathcal{M}_k(\mathbb{C})$ with respect to E. Then $(\mathcal{M}_k(\mathbb{C}), E) \underset{\mathcal{D}}{*} (\mathcal{M}_k(\mathbb{C}), E)$ is an MF algebra.

Theorem 3.4 Let $\mathcal{M}_k(\mathbb{C}) \supseteq \mathcal{A} \supseteq \mathcal{D} \subseteq \mathcal{B} \subseteq \mathcal{M}_k(\mathbb{C})$ be unital C*-inclusions of finite-dimensional C*-algebras and $E : \mathcal{M}_k(\mathbb{C}) \to \mathcal{D}$ be the trace preserving conditional expectation from $\mathcal{M}_k(\mathbb{C})$ onto $\mathcal{D}$. Then $(\mathcal{A}, E|_{\mathcal{A}}) *_{\mathcal{D}} (\mathcal{B}, E|_{\mathcal{B}})$ is MF.

The following is the main result of the paper.

Theorem 3.5 Suppose that $\mathcal{A}_1$ and $\mathcal{A}_2$ are two unital UHF-algebras with faithful tracial states $\tau_{\mathcal{A}_1}$ and $\tau_{\mathcal{A}_2}$ respectively. Let $\mathcal{A}_1 \supseteq \mathcal{D} \subseteq \mathcal{A}_2$ be unital inclusions of C*-algebras where $\mathcal{D}$ is a finite-dimensional C*-algebra. Assume that $E_{\mathcal{A}_1} : \mathcal{A}_1 \to \mathcal{D}$ and $E_{\mathcal{A}_2} : \mathcal{A}_2 \to \mathcal{D}$ are the trace preserving conditional expectations from $\mathcal{A}_1$ and $\mathcal{A}_2$ onto $\mathcal{D}$ respectively. Then the reduced amalgamated free product $(\mathcal{A}_1, E_{\mathcal{A}_1}) *_{\mathcal{D}} (\mathcal{A}_2, E_{\mathcal{A}_2})$ is an MF algebra if and only if $\tau_{\mathcal{A}_1}(z) = \tau_{\mathcal{A}_2}(z)$ for all $z \in \mathcal{D}$.

A brief overview of this paper is as follows. In Section 2, we recall the definitions of MF algebras and reduced amalgamated free product of unital C*-algebras Section 3 is devoted to results on reduced amalgamated free products of C*-algebras. We first show that a reduced free product of full matrix algebras with amalgamation over a full matrix subalgebra is MF. Later, we obtain some results about reduced amalgamated free products of two finite-dimensional C*-algebras with amalgamation over finite-dimensional C*-algebras. Finally, we give a necessary and sufficient condition for a reduced free product of two UHF algebras with amalgamated over a common finite-dimensional C*-subalgebra with respect to trace preserving conditional expectations to be MF.

2 Definitions and Preliminaries

2.1 Definition of MF Algebras

Suppose $\{\mathcal{M}_{k_n}(\mathbb{C})\}_{n=1}^{\infty}$ is a sequence of complex matrix algebras. We introduce the C*-direct product $\prod_{m=1}^{\infty}\mathcal{M}_{k_m}(\mathbb{C})$ of $\{\mathcal{M}_{k_n}(\mathbb{C})\}_{n=1}^{\infty}$ as follows:

$$\prod_{n=1}^{\infty}\mathcal{M}_{k_n}(\mathbb{C})=\{(Y_n)_{n=1}^{\infty} \mid \forall\, n\geq 1,\ Y_n\in\mathcal{M}_{k_n}(C)\ \text{ and }\ \|(Y_n)_{n=1}^{\infty}\|=\sup_{n\geq 1}\|Y_n\|<\infty\}.$$

Furthermore, we can introduce a norm-closed two sided ideal in $\prod_{n=1}^{\infty}\mathcal{M}_{k_n}(\mathbb{C})$ as follows:

$$\sum_{n=1}^{\infty}\mathcal{M}_{k_n}(\mathbb{C})=\left\{(Y_n)_{n=1}^{\infty}\in\prod_{n=1}^{\infty}\mathcal{M}_{k_n}(\mathbb{C}):\lim_{n\to\infty}\|Y_n\|=0\right\}.$$

Let π be the quotient map from $\prod_{n=1}^{\infty}\mathcal{M}_{k_n}(\mathbb{C})$ to $\prod_{n=1}^{\infty}\mathcal{M}_{k_n}(\mathbb{C})/\sum\limits_{n=1}^{\infty}\mathcal{M}_{k_n}(\mathbb{C})$. Then

$$\prod_{n=1}^{\infty}\mathcal{M}_{k_n}(\mathbb{C})/\sum_{n=1}^{\infty}\mathcal{M}_{k_n}(\mathbb{C})$$

is a unital C*-algebra. If we denote $\pi\left((Y_n)_{n=1}^{\infty}\right)$ by $[(Y_n)_n]$, then

$$\|[(Y_n)_n]\|=\limsup_{n\to\infty}\|Y_n\|\,.$$

Now we are ready to recall an equivalent definition of MF algebras given by Blackadar and Kirchberg [4].

Definition 2.1 *(Theorem 3.2.2, [4]) Let $\mathcal{A}$ be a separable C*-algebra. If $\mathcal{A}$ can be embedded as a C*-subalgebra of $\prod_{n=1}^{\infty}\mathcal{M}_{k_n}(\mathbb{C})/\sum\limits_{n=1}^{\infty}\mathcal{M}_{k_n}(\mathbb{C})$ for a sequence $\{k_n\}_{n=1}^{\infty}$ of integers, then $\mathcal{A}$ has MF property.*

The examples of MF algebras contain all finite dimensional C*-algebras, AF (approximately finite dimensional) algebras and quasidiagonal C*-algebras. In [13], Haagerup and Thorbjϕrnsen showed that $C_r^*(F_n)$ is an MF algebra for $n\geq 2$. For more examples of MF algebras, we refer the reader to [4] and [15].

2.2 Definition of Reduced Amalgamated Free Product of C*-algebras

In this section, we will explain the KSGNS representation of unital conditional expectations and recall a characterization of reduced amalgamated free product of C*-algebras. First, we give the definition of conditional expectation as follows:

Definition 2.2 *Let $\mathcal{D}$ be a unital C*-subalgebra of a unital C*-algebra $\mathcal{A}$. A projection from $\mathcal{A}$ onto $\mathcal{D}$ is a linear map $E : \mathcal{A} \to \mathcal{D}$ such that $E(d) = d$ for every $d \in \mathcal{D}$. A conditional expectation from $\mathcal{A}$ onto $\mathcal{D}$ is a contractive completely positive projection E from $\mathcal{A}$ onto $\mathcal{D}$ such that $E(dxd') = dE(x)d'$ for every $x \in \mathcal{A}$ and $d, d' \in \mathcal{D}$ (i.e. E is a $\mathcal{D}$-bimodule map).*

Now, we present a summary of the KSGNS representation of C*-algebras. Let $\mathcal{D}$ be a unital C*-algebra and let $\mathcal{A}$ be a unital C*-algebra, which contains a copy of $\mathcal{D}$ as a unital C*-subalgebra. We also suppose that there is a conditional expectation $E : \mathcal{A} \to \mathcal{D}$, satisfying

$$\forall a \in \mathcal{A} \text{ with } a \neq 0,\ \exists x \in \mathcal{A} \text{ such that } E(x^* a^* a x) \neq 0 \qquad (1).$$

The conditional expectations which satisfy (1) are called non-degenerate. Let $\mathcal{H} = L^2(\mathcal{A}, E)$ be the right Hilbert $\mathcal{D}$-module obtained from $\mathcal{A}$ by separation and completion with respect to the norm $\|a\| = \|\langle a, a\rangle_{\mathcal{H}}\|^{1/2}$, where $\langle a_1, a_2\rangle_{\mathcal{H}} = E(a_1^* a_2)$. Then the linear space $\mathcal{L}(\mathcal{H})$ of all adjointable $\mathcal{D}$-module operators on $\mathcal{H}$ is actually a C*-algebra and we have a representation $\pi : \mathcal{A} \to \mathcal{L}(\mathcal{H})$ defined by $\pi(a)\widehat{a'} = \widehat{aa'}$, where by $\widehat{a}$ we denote the element of $\mathcal{H}$, corresponding to $a \in \mathcal{A}$. π is faithful if E satisfies condition (1). In this construction, we have the specified element $\xi = \widehat{I_A} \in \mathcal{H}$. We call the triple $(\pi, \mathcal{H}, \xi)$ the KSGNS representation of $(\mathcal{A}, E)$. If $\mathcal{D} = \mathbb{C}$, then the Hilbert C*–module considered above become simply Hilbert space.

Let I be an index set. Card$(I) \geq 2$. Let $\mathcal{D}$ be a unital C*-algebra, and for each $i \in I$, we have a unital C*-algebra $\mathcal{A}_i$, which contains a copy of $\mathcal{D}$ as a unital C*-subalgebra. We also suppose that, for each $i \in I$, $E_i : \mathcal{A}_i \to \mathcal{D}$, is non-degenerate. Let $(\pi_i, \mathcal{H}_i, \xi_i)$ denote the KSGNS representations of $(\mathcal{A}_i, E_i)$. Note that each π_i is faithful. The Hilbert $\mathcal{D}$-submodule $\xi_i\mathcal{D} \subseteq \mathcal{H}_i$ is isomorphic to $\mathcal{D}$ and $\mathcal{H}_i \cong \xi_i \oplus \mathcal{H}_i^\circ$. Recall that $\mathcal{A}_i^\circ = \mathcal{A}_i \cap \ker E_i$ satisfies $\mathcal{D}\mathcal{A}_i^\circ\mathcal{D} \subseteq \mathcal{A}_i^\circ$ and $\mathcal{H}_i^\circ = \overline{\mathcal{A}_i^\circ \xi_i}$ is a C*-correspondence over $\mathcal{D}$. We define the free product Hilbert $\mathcal{D}$-module $(\mathcal{H}, \xi) = *(\mathcal{H}_i, \xi_i)$ by

$$\mathcal{H} = \xi\mathcal{D} \oplus \bigoplus_{n \geq 1} \bigoplus_{i_1 \neq \cdots \neq i_n} \mathcal{H}_{i_1}^\circ \otimes_{\mathcal{D}} \cdots \otimes_D \mathcal{H}_{i_n}^\circ.$$

Here, $\xi\mathcal{D}$ is the trivial Hilbert $\mathcal{D}$-module $\mathcal{D}$ with $\xi = \widehat{1}$ and $\otimes_{\mathcal{D}}$ means interior tensor product (see Lance's book [21]). For each $i \in I$, let

$$\mathcal{H}(i) = \xi\mathcal{D} \oplus \bigoplus_{n \geq 1} \bigoplus_{i \neq i_1 \neq \cdots \neq i_n} \mathcal{H}_{i_1}^\circ \otimes_{\mathcal{D}} \cdots \otimes_D \mathcal{H}_{i_n}^\circ$$

and define an isomorphism $U_i \in \mathcal{L}(\mathcal{H}_i \otimes_{\mathcal{D}} \mathcal{H}(i), \mathcal{H})$ by

$$U_i : \begin{array}{c} \xi_i\mathcal{D} \otimes_{\mathcal{D}} \xi\mathcal{D} \\ \mathcal{H}_i^\circ \otimes_{\mathcal{D}} \xi\mathcal{D} \\ \xi_i\mathcal{D} \otimes_{\mathcal{D}} \left(\mathcal{H}_{i_1}^\circ \otimes_{\mathcal{D}} \cdots \otimes_{\mathcal{D}} \mathcal{H}_{i_n}^\circ\right) \\ \mathcal{H}_i^\circ \otimes_{\mathcal{D}} \left(\mathcal{H}_{i_1}^\circ \otimes_{\mathcal{D}} \cdots \otimes_{\mathcal{D}} \mathcal{H}_{i_n}^\circ\right) \end{array} \xrightarrow{\cong} \begin{array}{c} \xi\mathcal{D} \\ \mathcal{H}_i^\circ \\ \mathcal{H}_{i_1}^\circ \otimes_{\mathcal{D}} \cdots \otimes_{\mathcal{D}} \mathcal{H}_{i_n}^\circ \\ \mathcal{H}_i^\circ \otimes_{\mathcal{D}} \mathcal{H}_{i_1}^\circ \otimes_{\mathcal{D}} \cdots \otimes_{\mathcal{D}} \mathcal{H}_{i_n}^\circ \end{array}$$

We define a *-representation $\lambda_i : \mathcal{A}_i \to \mathcal{L}(\mathcal{H})$ by $\lambda_i(x) = U_i(\pi_i \otimes 1)(x) U_i^*$. Observe that $\lambda_i|_{\mathcal{D}}$ is the canonical left action of $\mathcal{D}$ on the C*-correspondence $\mathcal{H}$ over $\mathcal{D}$, and hence $\lambda_i|_{\mathcal{D}} = \lambda_j|_{\mathcal{D}}$ for every i and j.

Definition 2.3 *The reduced amalgamated free product* $(\mathcal{A}, E) = *_{\mathcal{D}}(\mathcal{A}_i, E_i)$ *is the C*-subalgebra of* $\mathcal{L}(\mathcal{H})$ *generated by* $\cup_{i\in I}\lambda_i(\mathcal{A}_i)$.

The following lemma characterizes the reduced amalgamated free product of C*-algebras. It is one of the key ingredients in the proof of our main result.

Theorem 2.1 *(Theorem 7.2 [8]) Let* $I \in \mathcal{D} \subseteq \mathcal{A}_i$ *be unital C*-algebras with non-degenerate conditional expectations* E_i *from* $\mathcal{A}_i$ *onto* $\mathcal{D}$, *and let* $(\mathcal{A}, E) = *_{\mathcal{D}}(\mathcal{A}_i, E_i)$.

1. *There is an inclusion* $I \in \mathcal{D} \subseteq \mathcal{A}$ *and a non-degenerate conditional expectation* $E : \mathcal{A} \to \mathcal{D}$.
2. *We have inclusions* $\mathcal{D} \subseteq \mathcal{A}_i \subseteq \mathcal{A}$ *which are compatible on* $\mathcal{D}$, *and* $\mathcal{A}$ *is generated by* $\cup\mathcal{A}_i$ *as a C*-algebra.*
3. *We have* $E|_{\mathcal{A}_i} = E_i$ *for every* i, *and the C*-subalgebras* $\mathcal{A}_i$ *are free over* $\mathcal{D}$ *in* $(\mathcal{A}, E)$. *Namely,* $E(a_1 \cdots a_n) = 0$ *for every* $a_j \in \mathcal{A}_{i_j}^{\circ}$ *with* $i_1 \neq \cdots \neq i_n$.

Moreover, the above conditions uniquely characterize the reduced amalgamated free product $(\mathcal{A}, E)$.

3 Reduced Amalgamated Free Products Of Matrix Algebras

3.1 $\mathcal{D}$ is a full matrix algebra

In this subsection, we consider a reduced free product of two matrix algebras $\mathcal{A}$ and $\mathcal{B}$ with amalgamation over a C*-subalgebra $\mathcal{D}$ when $\mathcal{D}$ is a full matrix algebra. The following lemma is well-known.

Lemma 3.1 *(Theorem 2.1, [10]) Let* $\mathcal{A} \supseteq \mathcal{D} \subseteq \mathcal{B}$ *be unital inclusions of unital C*-algebras. Assume* $E_1 : \mathcal{A} \to \mathcal{D}$ *and* $E_2 : \mathcal{B} \to \mathcal{D}$ *are non-degenerate conditional expectations. Let* $\mathcal{M} = (\mathcal{A}, E_1) *_{\mathcal{D}} (\mathcal{B}, E_2)$ *be the reduced amalgamated free product of* $\mathcal{A}$ *and* $\mathcal{B}$ *with respect to* E_1 *and* E_2. *Suppose that there is a projection* $p \in \mathcal{D}$ *and there are partial isometries* $v_1, \cdots, v_n \in \mathcal{D}$ *such that* $v_i^* v_i \leq p$ *and* $\sum_{i=1}^n v_i v_i^* = 1 - p$. *Then*

$$p\mathcal{M}p = (p\mathcal{A}p, E_1|_{p\mathcal{A}p}) *_{p\mathcal{D}p} (p\mathcal{B}p, E_2|_{p\mathcal{B}p}).$$

By emulating the argument in the proof of Lemma 2.1 in [5], we obtain the following result.

Lemma 3.2 *Let $\mathcal{A}$ be a unital C*-algebra and suppose there is a projection $p \in \mathcal{A}$ and there are partial isometries $v_1, \cdots, v_n \in \mathcal{A}$ such that $v_i^* v_i \leq p$ and $\sum_{i=1}^{n} v_i v_i^* = 1 - p$. Then $\mathcal{A}$ is MF if and only if $p\mathcal{A}p$ is MF.*

The following lemma can be found in [16].

Lemma 3.3 *(Theorem 3.4.1, [16]) Suppose that $\{\mathcal{A}_i : i = 1, \cdots, n\}$ is a family of unital separable AH (approximately homogeneous) algebras with faithful tracial state τ_i for $i \in \{1, \cdots, n\}$. Then*

$$(\mathcal{A}_1, \tau_1) *_{red} \cdots *_{red} (\mathcal{A}_n, \tau_n)$$

is an MF algebra.

Now, we are able to obtain our first result on the reduced amalgamated free products of matrix algebras.

Theorem 3.1 *Suppose that $k, n, m \in \mathbb{N}$. If $\mathcal{M}_k(\mathbb{C})$ can be unitally embedded into $\mathcal{M}_n(\mathbb{C})$ and $\mathcal{M}_m(\mathbb{C})$ as a unital C*-subalgebra respectively, then for the trace preserving conditional expectations E_1 and E_2 from $\mathcal{M}_n(\mathbb{C})$ and $\mathcal{M}_m(\mathbb{C})$ onto $\mathcal{M}_k(\mathbb{C})$,*

$$(\mathcal{M}_n(\mathbb{C}), E_1) *_{\mathcal{M}_k(\mathbb{C})} (\mathcal{M}_m(\mathbb{C}), E_2)$$

is an MF algebra.

Proof. Since $\mathcal{M}_k(\mathbb{C})$ can be unitally embedded into $\mathcal{M}_n(\mathbb{C})$ and $\mathcal{M}_m(\mathbb{C})$ respectively, by Lemma 6.6.3 in [19], we have that

$$\mathcal{M}_n(\mathbb{C}) \cong \mathcal{M}_k(\mathbb{C}) \otimes \mathcal{M}_{\frac{n}{k}}(\mathbb{C})$$

and

$$\mathcal{M}_m(\mathbb{C}) \cong \mathcal{M}_k(\mathbb{C}) \otimes \mathcal{M}_{\frac{m}{k}}(\mathbb{C}).$$

This implies that, for a one-dimensional projection $p \in \mathcal{M}_k(\mathbb{C})$, we have

$$p\mathcal{M}_n(\mathbb{C})p \cong \mathcal{M}_{\frac{n}{k}}(\mathbb{C}) \text{ and } p\mathcal{M}_m(\mathbb{C})p \cong \mathcal{M}_{\frac{m}{k}}(\mathbb{C}).$$

Suppose τ_1 and τ_2 are the faithful tracial states on $\mathcal{M}_{\frac{n}{k}}(\mathbb{C})$ and $\mathcal{M}_{\frac{n}{k}}(\mathbb{C})$ respectively. Let $E_1 = id_k \otimes \tau_1$ and $E_2 = id_k \otimes \tau_2$ where id_k is the identity map on $\mathcal{M}_k(\mathbb{C})$. Then E_1 and E_2 are conditional expectations from $\mathcal{M}_k(\mathbb{C}) \otimes \mathcal{M}_{\frac{n}{k}}(\mathbb{C})$ and $\mathcal{M}_k(\mathbb{C}) \otimes \mathcal{M}_{\frac{m}{k}}(\mathbb{C})$ onto $\mathcal{M}_k(\mathbb{C})$ respectively. It is easy to check that E_1 and E_2 are trace preserving. Let

$$\mathcal{M} = (\mathcal{M}_n(\mathbb{C}), E_1) *_{\mathcal{M}_k(\mathbb{C})} (\mathcal{M}_m(\mathbb{C}), E_2).$$

Since there are partial isometries $v_1, \cdots, v_{k-1} \in \mathcal{M}_k(\mathbb{C})$ such that $v_i^* v_i = p$ for each $i \in \{1, \cdots, k-1\}$ and $\sum_{i=1}^{k-1} v_i v_i^* = I_{\mathcal{M}_k(\mathbb{C})} - p$, by Lemma 3.1, we have

$$\begin{aligned} p\mathcal{M}p &= \left(p\mathcal{M}_n(\mathbb{C})p, E_1|_{p\mathcal{M}_n(\mathbb{C})p}\right) *_{p\mathcal{M}_k(\mathbb{C})p} \left(p\mathcal{M}_m(\mathbb{C})p, E_2|_{p\mathcal{M}_m(\mathbb{C})p}\right) \\ &\cong \left(\mathcal{M}_{\frac{n}{k}}(\mathbb{C}), \tau_1\right) *_{red} \left(\mathcal{M}_{\frac{m}{k}}(\mathbb{C}), \tau_2\right) \qquad (3.1) \end{aligned}$$

Combining (3.1) with Lemma 3.3 and Lemma 3.2, we conclude that

$$\left(\mathcal{M}_n(\mathbb{C}), E_1\right) *_{\mathcal{M}_k(\mathbb{C})} \left(\mathcal{M}_m(\mathbb{C}), E_2\right)$$

is an MF algebra as desired. ■

3.2 $\mathcal{D}$ is a finite-dimensional abelian C*-algebra

More generally, we consider the case when the overlap C*-algebra $\mathcal{D}$ is a finite-dimensional C*-algebra. In this subsection we consider a reduced free product of $\mathcal{M}_k(\mathbb{C})$ and $\mathcal{M}_k(\mathbb{C})$ with amalgamation over a finite-dimensional abelian C*-subalgebra $\mathcal{D}$ with respect to a trace preserving conditional expectation. The following lemma can be found in [4].

Lemma 3.4 *(Proposition 3.3.6, [4]) If $\mathcal{A}$ and $\mathcal{B}$ are MF algebras and one of them is nuclear, then $\mathcal{A} \otimes \mathcal{B}$ is MF.*

It is well-known that the full matrix algebra $\mathcal{M}_n(\mathbb{C})$ and the reduced C*-algebra $C_r^*(F_n)$ of the free group F_n have canonical tracial states, without loss of generality, we denote the reduced free product of $\mathcal{M}_n(\mathbb{C})$ and $C_r^*(F_n)$ with respect to their canonical tracial states by $\mathcal{M}_n(\mathbb{C}) *_{red} C_r^*(F_n)$. Similarly, the reduced free product of $\mathcal{M}_n(\mathbb{C})$ and $\mathcal{M}_m(\mathbb{C})$ with respect to their tracial states is denoted by $\mathcal{M}_n(\mathbb{C}) *_{red} \mathcal{M}_m(\mathbb{C})$.

For the purpose of simplicity, we will first assume that $\mathcal{D}$ is a two dimensional abelian C*-subalgebra of $\mathcal{M}_k(\mathbb{C})$. Let $\mathcal{D} \subseteq \mathcal{M}_k(\mathbb{C})$ be a two-dimensional abelian C*-subalgebra of $\mathcal{M}_k(\mathbb{C})$, i.e., there are non-zero orthogonal projections P_1, P_2 in $\mathcal{M}_k(\mathbb{C})$ such that

$$\mathcal{D} = \mathbb{C}P_1 \oplus \mathbb{C}P_2.$$

Let $E : \mathcal{M}_k(\mathbb{C}) \to \mathcal{D}$ be a trace preserving conditional expectation from $\mathcal{M}_k(\mathbb{C})$ onto $\mathcal{D}$ and $\left(\mathcal{M}_k(\mathbb{C}), E\right) *_{\mathcal{D}} \left(\mathcal{M}_k(\mathbb{C}), E\right)$ be the reduced free product of $\left(\mathcal{M}_k(\mathbb{C}), E\right)$ and $\left(\mathcal{M}_k(\mathbb{C}), E\right)$ with amalgamation over $\mathcal{D}$ with respect to the conditional expectation E. By Corollary II.6.10.8 [3], E is the unique trace preserving conditional expectation from $\mathcal{M}_k(\mathbb{C})$ onto $\mathcal{D}$. Assume that $k_1 = \text{rank}(P_1)$ and $k_2 = \text{rank}(P_2)$. Consider C*-algebras $C_r^*(F_2)$,

$$\left(\mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})\right) \otimes \mathcal{M}_k(\mathbb{C})$$

and

$$\left(C_r^*(F_2) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})\right) \otimes \mathcal{M}_k(\mathbb{C}).$$

Then, by the definition of reduced free product of C*-algebras, $C_r^*(F_2) \otimes \mathcal{M}_k(\mathbb{C})$ and $\left(\mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})\right) \otimes \mathcal{M}_k(\mathbb{C})$ can be treated as unital C*-subalgebras of

$$\left(C_r^*(F_2) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})\right) \otimes \mathcal{M}_k(\mathbb{C}).$$

Let U_1, U_2 are canonical generators of $C_r^*(F_2)$. Then it is easy to see that the unitary

$$U = U_1 \otimes P_1 + U_2 \otimes P_2$$

is in

$$C_r^*(F_2) \otimes \mathcal{M}_k(\mathbb{C}) \subseteq \left(C_r^*(F_2) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})\right) \otimes \mathcal{M}_k(\mathbb{C}).$$

From above notations, we can get the following technical result.

Lemma 3.5 *Suppose $P_1, P_2 \in \mathcal{M}_k(\mathbb{C})$ are orthogonal projections from above. Then there is a C*-subalgebra $\mathcal{M}$ of*

$$\left(\mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})\right) \otimes \mathcal{M}_k(\mathbb{C}) \subseteq \left(C_r^*(F_2) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})\right) \otimes \mathcal{M}_k(\mathbb{C})$$

such that:

1. $\mathcal{M} \cong \mathcal{M}_k(\mathbb{C})$,
2. $\mathcal{M}_{k_1}(\mathbb{C}) \otimes P_1 + \mathcal{M}_{k_2}(\mathbb{C}) \otimes P_2 \subseteq \mathcal{M}$.

Proof. Let τ_1 be the faithful tracial state on $\mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})$ obtained from faithful tracial states on $\mathcal{M}_{k_1}(\mathbb{C})$ and $\mathcal{M}_{k_2}(\mathbb{C})$, and τ_2 be the unique faithful tracial state on $\mathcal{M}_k(\mathbb{C})$. Then the state $\tau = \tau_1 \otimes \tau_2$ is a faithful tracial state on $\left(\mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})\right) \otimes \mathcal{M}_k(\mathbb{C})$. Let $\{e_{ij}\}_{i,j=1}^{k_1}$, $\{g_{ij}\}_{i,j=1}^{k_2}$ and $\{f_{ij}\}_{i,j=1}^{k}$ be systems of matrix units of $\mathcal{M}_{k_1}(\mathbb{C})$, $\mathcal{M}_{k_2}(\mathbb{C})$ and $\mathcal{M}_k(\mathbb{C})$, respectively. So we have

$$\begin{aligned} \tau(e_{11} \otimes P_1) &= \frac{1}{k_1} \cdot \frac{k_1}{k} = 1 \cdot \frac{1}{k} \\ &= \tau(I \otimes f_{11}) = \frac{1}{k_2} \cdot \frac{k_2}{k} \\ &= \tau(g_{11} \otimes P_2) \end{aligned}$$

where I is the unit in $\mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})$. Note that $\mathcal{M}_{k_1}(\mathbb{C}) \otimes \mathcal{M}_k(\mathbb{C})$ is *-isomorphic to a full matrix algebra. Thus there is a partial isometry W_1 in $\mathcal{M}_{k_1}(\mathbb{C}) \otimes \mathcal{M}_k(\mathbb{C})$ such that

$$W_1 W_1^* = e_{11} \otimes P_1; \; W_1^* W_1 = I \otimes f_{11}.$$

Similarly, there is a partial isometry $W_2 \in \mathcal{M}_{k_2}(\mathbb{C}) \otimes \mathcal{M}_k(\mathbb{C})$ such that

$$W_2 W_2^* = g_{11} \otimes P_2; \; W_1^* W_1 = I \otimes f_{11}.$$

Let $V_1 = W_2^* W_1$. Then V_1 is a partial isometry in $\left(\mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})\right) \otimes \mathcal{M}_k$ satisfying

$$V_1^* V_1 = e_{11} \otimes P_1 \text{ and } V_1 V_1^* = g_{11} \otimes P_2.$$

Therefore the C*-subalgebra $\mathcal{M}$ in

$$\left(\mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})\right) \otimes \mathcal{M}_k(\mathbb{C})$$

generated by $\{e_{ij}\}_{i,j=1}^{k_1} \otimes P_1$, $\{g_{ij}\}_{i,j=1}^{k_2} \otimes P_2$ and V_1 is *-isomorphic to the full matrix algebra $\mathcal{M}_k(\mathbb{C})$. It is easy to see that

$$\mathcal{M}_{k_1}(\mathbb{C}) \otimes P_1 + \mathcal{M}_{k_2}(\mathbb{C}) \otimes P_2 \subseteq \mathcal{M}.$$

■

Let $C^*(\mathcal{M}, U\mathcal{M}U^*)$ be the C*-algebra generated by $\mathcal{M}$ and $U\mathcal{M}U^*$ in

$$\left(C_r^*(F_2) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})\right) \otimes \mathcal{M}_k(\mathbb{C}).$$

Since

$$C_r^*(F_2) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C}) \cong C_r^*(\mathbb{Z}) *_{red} C_r^*(\mathbb{Z}) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C}),$$

we know that

$$\left(C_r^*(F_2) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})\right) \otimes \mathcal{M}_k(\mathbb{C})$$

is an MF algebra by Lemma 3.4 and 3.3. This implies that $C^*(\mathcal{M}, U\mathcal{M}U^*)$ is an MF algebra. Notice that

$$U(I \otimes P_1)U^* = U_1 I U_1^* \otimes P_1 = I \otimes P_1$$

and

$$U(I \otimes P_2)U^* = U_2 I U_2^* \otimes P_2 = I \otimes P_2,$$

this implies that

$$U\left[\mathbb{C}(I \otimes P_1) \oplus \mathbb{C}(I \otimes P_2)\right]U^* = \mathbb{C}(I \otimes P_1) \oplus \mathbb{C}(I \otimes P_2).$$

Therefore $\mathbb{C}(I \otimes P_1) \oplus \mathbb{C}(I \otimes P_2) \subseteq U\mathcal{M}U^*$.

Now, we consider the reduced amalgamated free product

$$\left(\mathcal{M}_k(\mathbb{C}), E\right) \underset{\mathcal{D}}{*} \left(\mathcal{M}_k(\mathbb{C}), E\right)$$

of $(\mathcal{M}_k(\mathbb{C}), E)$ and $(\mathcal{M}_k(\mathbb{C}), E)$ when $\mathcal{D}$ is a two-dimensional abelian C*-algebra.

Proposition 3.1 *Let $\mathcal{D} \subseteq \mathcal{M}_k(\mathbb{C})$ be a unital inclusion of C*-algebras where $\mathcal{D}$ is a two-dimensional abelian C*-algebra. Suppose $E : \mathcal{M}_k(\mathbb{C}) \to \mathcal{D}$ is the trace preserving conditional expectation from $\mathcal{M}_k(\mathbb{C})$ onto $\mathcal{D}$. Let U and $\mathcal{M}$ be as above, then*

$$C^*(\mathcal{M}, U\mathcal{M}U^*) \cong \left(\mathcal{M}_k(\mathbb{C}), E\right) \underset{\mathcal{D}}{*} \left(\mathcal{M}_k(\mathbb{C}), E\right).$$

Therefore $\left(\mathcal{M}_k(\mathbb{C}), E\right) \underset{\mathcal{D}}{*} \left(\mathcal{M}_k(\mathbb{C}), E\right)$ is an MF algebra.

Proof. Let τ_1 be the faithful tracial state on $C_r^*(F_2) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})$ obtained from tracial states on $C_r^*(F_2)$, $\mathcal{M}_{k_1}(\mathbb{C})$ and $\mathcal{M}_{k_2}(\mathbb{C})$, and τ_2 be the faithful tracial state on $\mathcal{M}_k(\mathbb{C})$. Then the state $\tau = \tau_1 \otimes \tau_2$ is a faithful tracial state on

$$(C_r^*(F_2) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})) \otimes \mathcal{M}_k(\mathbb{C}).$$

Let I be the unit in $C_r^*(F_2) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})$. It is clear that $\mathbb{C}(I \otimes P_1) \oplus \mathbb{C}(I \otimes P_2)$ is *-isomorphic to $\mathcal{D}$. Assume $\mathcal{M}_1 = \mathcal{M}$ and $\mathcal{M}_2 = U\mathcal{M}U^*$. From the above discussion, we have that

$$\mathbb{C}(I \otimes P_1) \oplus \mathbb{C}(I \otimes P_2) \subseteq \mathcal{M}_1 \text{ and } \mathbb{C}(I \otimes P_1) \oplus \mathbb{C}(I \otimes P_2) \subseteq \mathcal{M}_2.$$

Define a linear mapping

$$\widetilde{E} : C^*(\mathcal{M}_1, \mathcal{M}_2) \to \mathbb{C}(I \otimes P_1) \oplus \mathbb{C}(I \otimes P_2)$$

by

$$\widetilde{E}(A) = \sum_{i=1}^{2} \frac{1}{\tau(I \otimes P_i)} (\tau(A(I \otimes P_i)) I) \otimes P_i$$

for $A \in C^*(\mathcal{M}, U\mathcal{M}U^*)$. By the fact that

$$\begin{aligned} & I \otimes P_1 \oplus I \otimes P_2 \\ = \;& \widetilde{I} \in (C_r^*(F_2) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})) \otimes \mathcal{M}_k(\mathbb{C}), \end{aligned}$$

it is not hard to see that $\widetilde{E}$ is a trace preserving conditional expectation. The restrictions of $\widetilde{E}$ on $\mathcal{M}_1$ and $\mathcal{M}_2$, i.e., $\widetilde{E}|_{\mathcal{M}_1}$ and $\widetilde{E}|_{\mathcal{M}_2}$ are trace preserving conditional expectations from $\mathcal{M}_1$ and $\mathcal{M}_2$ onto $\mathbb{C}(I \otimes P_1) \oplus \mathbb{C}(I \otimes P_2)$ respectively. Since $\mathcal{M}_1 \cong \mathcal{M}_k(\mathbb{C})$ and the trace preserving conditional expectation from $\mathcal{M}_k(\mathbb{C})$ onto $\mathcal{D}$ is unique, for showing $C^*(\mathcal{M}_1, \mathcal{M}_2)$ is *-isomorphic to $(\mathcal{M}_k(\mathbb{C}), E) *_{\mathcal{D}} (\mathcal{M}_k(\mathbb{C}), E)$, we only need to show that

$$\widetilde{E}(T_1 T_2 \cdots T_n) = 0$$

if $T_j \in \mathcal{M}_{l_j}$, $\widetilde{E}(T_j) = 0$ for any $j \in \{1, \cdots, n\}$ and $l_1 \neq l_2 \neq \cdots \neq l_n$, $l_1, \cdots, l_n \in \{1, 2\}$ by Theorem 2.1. Suppose X_i, $Y_i \in \mathcal{M}_1$ for $i \in \{1, \cdots, n\}$ with $\widetilde{E}(X_i) = \widetilde{E}(Y_i) = 0$. If we can show that

$$\widetilde{E}(UX_1U^*Y_1UX_2U^*Y_2 \cdots UX_nU^*Y_n) = 0$$

for the reduced word $UX_1U^*Y_1UX_2U^*Y_2 \cdots UX_nU^*Y_n$ in $C^*(\mathcal{M}_1, \mathcal{M}_2)$, then, for the other forms of reduced word, the desired result will be followed by using similar arguments. Therefore we can conclude that $(\mathcal{M}_k(\mathbb{C}), E) *_{\mathcal{D}} (\mathcal{M}_k(\mathbb{C}), E) \cong C^*(\mathcal{M}, U\mathcal{M}U^*)$. It follows that $(\mathcal{M}_k(\mathbb{C}), E) *_{\mathcal{D}} (\mathcal{M}_k(\mathbb{C}), E)$ is an MF algebra.

Suppose $X_i = X_{11}^{(i)} + X_{22}^{(i)} + X_{12}^{(i)} + X_{21}^{(i)}$ with

$$X_{11}^{(i)} = x_{11}^{(i)} \otimes P_1 = \sum_{s_1=1, t=1}^{s_1=k_1, t=k_1} x_{s_1 t}^{(i)} \otimes f_{s_1 t}$$

where $x_{11}^{(i)} \in \mathcal{M}_{k_1}(\mathbb{C})$ and $x_{s_1 s_1}^{(i)} = x_{11}^{(i)}$, $x_{s_1 t}^{(i)} = 0$ if $s_1 \neq t$, and

$$X_{22}^{(i)} = x_{22}^{(i)} \otimes P_2 = \sum_{s_2=k_1+1, h=k_1+1}^{s_2=k, h=k} x_{s_2 h}^{(i)} \otimes f_{s_2 h}$$

where $x_{s_2 s_2}^{(i)} = x_{22}^{(i)} \in \mathcal{M}_{k_2}(\mathbb{C})$, $x_{s_2 h}^{(i)} = 0$ if $s_2 \neq h$, and

$$X_{12}^{(i)} = \sum_{s_1=1, s_2=k_1+1}^{s_1=k_1, s_2=k} x_{s_1 s_2}^{(i)} \otimes f_{s_1 s_2} \quad \text{where } x_{s_1 s_2}^{(i)} \in \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})$$

as well as

$$X_{21}^{(i)} = \sum_{s_1=1, s_2=k_1+1}^{s_1=k_1, s_2=k} x_{s_2 s_1}^{(i)} \otimes f_{s_2 s_1} \text{ where } x_{s_2 s_1}^{(i)} \in \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C}).$$

Similarly, suppose $Y_i = Y_{11}^{(i)} + Y_{22}^{(i)} + Y_{12}^{(i)} + Y_{21}^{(i)}$ with

$$Y_{11}^{(i)} = y_{11}^{(i)} \otimes P_1 = \sum_{s_1=1, t=1}^{s_1=k_1, t=k_1} y_{s_1 t}^{(i)} \otimes f_{s_1 t}$$

where $y_{s_1 s_1}^{(i)} = y_{11}^{(i)} \in \mathcal{M}_{k_1}(\mathbb{C})$, $y_{s_1 t}^{(i)} = 0$ if $s_1 \neq t$, and

$$Y_{22}^{(i)} = y_{22}^{(i)} \otimes P_2 = \sum_{s_2=k_1+1, h=k_1+1}^{s_2=k, h=k} y_{s_2 h}^{(i)} \otimes f_{s_2 h}$$

where $y_{s_2 s_2}^{(i)} = y_{22}^{(i)} \in \mathcal{M}_{k_2}(\mathbb{C})$, $y_{s_2 h}^{(i)} = 0$ if $s_2 \neq h$, and

$$Y_{12}^{(i)} = \sum_{s_1=1, s_2=k_1+1}^{s_1=k_1, s_2=k} y_{s_1 s_2}^{(i)} \otimes f_{s_1 s_2} \text{ where } y_{s_1 s_2}^{(i)} \in \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C})$$

as well as

$$Y_{21}^{(i)} = \sum_{s_1=1, s_2=k_1+1}^{s_1=k_1, s_2=k} y_{s_2 s_1}^{(i)} \otimes f_{s_2 s_1} \text{ where } y_{s_2 s_1}^{(i)} \in \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C}).$$

Note that

$$(I \otimes P_1) X_i (I \otimes P_1) = X_{11}^{(i)}, \quad (I \otimes P_2) X_i (I \otimes P_2) = X_{22}^{(i)}.$$

Since $\widetilde{E}(X_i) = 0$, we have

$$\tau\left(x_{11}^{(i)} \otimes P_1\right) = \tau\left(x_{22}^{(i)} \otimes P_1\right) = 0$$

for each $i \in \{1, \cdots, n\}$. Similarly,

$$\tau\left(y_{11}^{(i)} \otimes P_2\right) = \tau\left(y_{22}^{(i)} \otimes P_2\right) = 0$$

for each $i \in \{1, \cdots, n\}$. This implies that

$$\tau_1\left(x_{11}^{(i)}\right) = \tau_1\left(x_{22}^{(i)}\right) = \tau_1\left(y_{11}^{(i)}\right) = \tau_1\left(y_{22}^{(i)}\right) = 0$$

for each $i \in \{1, \cdots, n\}$. Therefore

$$\tau_1\left(x_{s_1 s_1}^{(i)}\right) = \tau_1\left(x_{s_2 s_2}^{(i)}\right) = \tau_1\left(y_{s_1 s_1}^{(i)}\right) = \tau_1\left(y_{s_2 s_2}^{(i)}\right) = 0 \tag{3.2}$$

for each $s_1 \in \{1, \cdots, k_1\}$ and each $s_2 \in \{k_1 + 1, \cdots, k\}$. Assume

$$UX_1U^*Y_1UX_2U^*Y_2 \cdots UX_nU^*Y_n = A_{11} + A_{22} + A_{12} + A_{21}$$

where $A_{ij} \in (I \otimes P_i)\, C^*(\mathcal{M}, U\mathcal{M}U^*)\,(I \otimes P_j)$ for $i, j \in \{1, 2\}$. Then

$$\begin{aligned} &A_{11} \\ &= \sum_{i_1, \cdots, i_{n-1}, j_1, \cdots, j_n \in \{1,2\}} (U_1 \otimes P_1)\, X_{1j_1}^{(1)} \left(U_{j_1}^* \otimes P_{j_1}\right) Y_{j_1 i_1}^{(1)} \cdots X_{i_{n-1} j_n}^{(n)} \left(U_{j_n}^* \otimes P_{j_n}\right) Y_{j_n 1} \end{aligned} \tag{3.3}$$

and

$$\begin{aligned} &A_{22} \\ &= \sum_{i_1, \cdots, i_{n-1}, j_1, \cdots, j_n \in \{1,2\}} (U_2 \otimes P_2)\, X_{2j_1}^{(1)} \left(U_{j_1}^* \otimes P_{j_1}\right) Y_{j_1 i_1} \cdots X_{i_{n-1} j_n}^{(n)} \left(U_{j_n}^* \otimes P_{j_n}\right) Y_{j_n 2}. \end{aligned} \tag{3.4}$$

So, for showing $\widetilde{E}\left(UX_1U^*Y_1UX_2U^*Y_2 \cdots UX_nU^*Y_n\right) = 0$, we only need to prove the following claims.

Claim 3.1 For every s_1 and every m in $\{1, \cdots, k_1\}$, every $s_2, l \in \{k_1 + 1, \cdots, k\}$ and $i_1, \cdots, i_{n-1}, j_1, \cdots, j_n \in \{1, 2\}$, we have

$$\tau_1\left(U_1 x_{m s_{j_1}}^{(1)} U_{j_1}^* y_{s_{j_1} s_{i_1}}^{(1)} \cdots U_{i_{n-1}} x_{s_{i_{n-1}} s_{j_n}}^{(n)} U_{j_n}^* y_{s_{j_n} m}^{(n)}\right) = 0$$

and

$$\tau_1\left(U_2 x_{l s_{j_1}}^{(1)} U_{j_1}^* y_{s_{j_1} s_{i_1}}^{(1)} \cdots U_{i_{n-1}} x_{s_{i_{n-1}} s_{j_n}}^{(n)} U_{j_n}^* y_{s_{j_n} l}^{(n)}\right) = 0$$

Proof For word $U_1 x_{m s_{j_1}}^{(1)} U_{j_1}^* y_{s_{j_1} s_{i_1}}^{(1)} \cdots U_{i_{n-1}} x_{s_{i_{n-1}} s_{j_n}}^{(n)} U_{j_n}^* y_{s_{j_n} m}^{(n)}$, let k denote the total number of elements in

$$\left\{x_{m s_{j_1}}^{(1)}, \cdots, x_{s_{i_{n-1}} s_{j_n}}^{(n)}, y_{s_{j_1} s_{i_1}}^{(1)}, \cdots, y_{s_{j_n} s_m}^{(n)}\right\} \subseteq \mathcal{M}_{k_1} * \mathcal{M}_{k_2}$$

which have non-zero trace value. If $k = 0$, then each element in

$$\left\{ x^{(1)}_{ms_{j_1}}, \cdots, x^{(n)}_{s_{i_{n-1}}s_{j_n}}, y^{(1)}_{s_{j_1}s_{i_1}}, \cdots, y^{(n)}_{s_{j_n}s_m} \right\},$$

has zero trace value. Since U_1 and U_2 are two free Haar unitaries in $C^*_r(F_2)$ and

$$U_1 x^{(1)}_{ms_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1}s_{i_1}} \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}}s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n}m}$$

is a reduced word in $C^*_r(F_2) *_{red} (\mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C}))$, we have

$$\tau_1 \left(U_1 x^{(1)}_{ms_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1}s_{i_1}} \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}}s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n}m} \right) = 0.$$

Assume that $k = 1$. Then there is an integer t such that $\tau_1\left(x^{(t)}_{s_{i_{t-1}}s_{j_t}}\right)$ or $\tau_1\left(y^{(t)}_{s_{j_t}s_{i_t}}\right)$ is non-zero. From equation (3.2), we get that $i_{t-1} \neq j_t$ or $j_t \neq i_t$. Without loss of generality, we may assume that $\tau_1\left(y^{(t)}_{s_{j_t}s_{i_t}}\right) \neq 0$, $1 = j_t \neq i_t = 2$ and let $\overset{\circ}{y}^{(t)}_{s_1s_2} = y^{(t)}_{s_1s_2} - \tau_1\left(y^{(t)}_{s_1s_2}\right)$. Therefore

$$\begin{aligned} & U_1 x^{(1)}_{ms_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1}s_{i_1}} \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}}s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n}m} \\ & = U_1 x^{(1)}_{ms_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1}s_{i_1}} \cdots U^*_1 \overset{\circ}{y}^{(t)}_{s_1s_2} U_2 \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}}s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n}m} \\ & + \tau_1\left(y^{(t)}_{s_1s_2}\right) U_1 x^{(1)}_{ms_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1}s_{i_1}} \cdots U^*_1 U_2 \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}}s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n}s_m} \end{aligned}$$

It is easy to see that

$$U_1 x^{(1)}_{m,s_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1}s_{i_1}} \cdots U^*_1 \overset{\circ}{y}^{(t)}_{s_1s_2} U_2 \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}}s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n}m}$$

and

$$U_1 x^{(1)}_{m,s_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1}s_{i_1}} \cdots U^*_1 U_2 \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}}s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n}s_m}$$

are reduced words in $C^*_r(F_2) *_{red} (\mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C}))$. Since $k = 1$, we have the total number of elements in each reduced word from above which belong to $\mathcal{M}_{k_1} * \mathcal{M}_{k_2}$ with non-zero trace is 0. It follows that

$$\begin{aligned} & \tau_1 \left(U_1 x^{(1)}_{ms_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1}s_{i_1}} \cdots U^*_1 \overset{\circ}{y}^{(t)}_{s_1s_2} U_2 \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}}s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n}m} \right) \\ = \; & \tau_1 \left(U_1 x^{(1)}_{ms_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1}s_{i_1}} \cdots U^*_1 U_2 \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}}s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n}m} \right) = 0. \end{aligned}$$

By a similar argument, if $k \geq 2$, we can decompose

$$U_1 x^{(1)}_{ms_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1}s_{i_1}} \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}}s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n}m}$$

as a sum of two reduced words in

$$C^*_r(F_2) *_{red} (\mathcal{M}_{k_1}(\mathbb{C}) *_{red} \mathcal{M}_{k_2}(\mathbb{C}))$$

such that, for each reduced word, the total number of elements in $\mathcal{M}_{k_1} * \mathcal{M}_{k_2}$ with non-zero trace is $k-1$. This implies that we can decompose

$$U_1 x^{(1)}_{m s_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1} s_{i_1}} \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}} s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n} m}$$

as a sum of finitely many reduced words satisfying the condition that the total number of elements which belong to $\mathcal{M}_{k_1} * \mathcal{M}_{k_2}$ in the reduced word with non-zero trace is 0. Therefore we have that the trace value of each summand is 0. It implies that

$$\tau_1 \left(U_1 x^{(1)}_{m s_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1} s_{i_1}} \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}} s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n} m} \right) = 0.$$

By the same argument, we conclude that

$$\tau_1 \left(U_2 x^{(1)}_{l s_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1} s_{i_1}} \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}} s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n} l} \right) = 0$$

So this complete the proof of Claim 3.1.

Claim 3.2 If

$$\tau_1 \left(U_1 x^{(1)}_{m s_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1} s_{i_1}} \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}} s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n} m} \right) = 0$$

and

$$\tau_1 \left(U_2 x^{(1)}_{l s_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1} s_{i_1}} \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}} s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n} l} \right) = 0$$

for each $s_1, m \in \{1, \cdots, k_1\}$, $s_2, l \in \{k_1+1, \cdots, k\}$ and $i_1, \cdots, i_{n-1}, j_1, \cdots, j_n \in \{1, 2\}$, then

$$\widetilde{E} \left(U X_1 U^* Y_1 U X_2 U^* Y_2 \cdots U X_n U^* Y_n \right) = 0.$$

Proof Suppose

$$\begin{aligned} & \left(U_1 \otimes P_1 \right) X^{(1)}_{1 j_1} \left(U^*_{j_1} \otimes P_{j_1} \right) Y^{(1)}_{j_1 i_1} \cdots \left(U_{i_{n-1}} \otimes P_{i_{n-1}} \right) X^{(n)}_{i_{n-1} j_n} \left(U^*_{j_n} \otimes P_{j_n} \right) Y^{(n)}_{j_n 1} \\ = & \sum_{s,t=1}^{k_1} T_{st} \otimes f_{st}. \end{aligned}$$

in (3.3). Then

$$T_{mm} = \sum_{s_2=k_1+1}^{k} \sum_{s_1=1}^{k_1} \sum_{j_1,\cdots,j_n,i_1,\cdots,i_{n-1} \in \{1,2\}} U_1 x^{(1)}_{m s_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1} s_{i_1}} \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}} s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n} m} \tag{3.5}$$

for $m \in \{1, \cdots, k_1\}$. Since

$$\begin{aligned} & \tau \left(\left(U_1 \otimes P_1 \right) X^{(1)}_{1 j_1} \left(U^*_{j_1} \otimes P_{j_1} \right) Y^{(1)}_{j_1 i_1} \cdots \left(U_{i_{n-1}} \otimes P_{i_{n-1}} \right) X^{(n)}_{i_{n-1} j_n} \left(U^*_{j_n} \otimes P_{j_n} \right) Y^{(n)}_{j_n 1} \right) \\ = & \sum_{m=1}^{k_1} \frac{1}{k} \tau_1 \left(T_{mm} \right) \end{aligned} \tag{3.6}$$

and

$$\tau_1\left(U_1 x^{(1)}_{ms_{j_1}} U^*_{j_1} y^{(1)}_{s_{j_1} s_{i_1}} \cdots U_{i_{n-1}} x^{(n)}_{s_{i_{n-1}} s_{j_n}} U^*_{j_n} y^{(n)}_{s_{j_n} m}\right) = 0$$

for each $s_1, m \in \{1, \cdots, k_1\}$, $s_2 \in \{k_1+1, \cdots, k\}$ and $i_{1,} \cdots, i_{n-1}, j_{1,} \cdots, j_n \in \{1,2\}$, from (3.5) and (3.6), we get

$$\tau\left(\left(U_1 \otimes P_1\right) X^{(1)}_{1j_1}\left(U^*_{j_1} \otimes P_{j_1}\right) Y^{(1)}_{j_1 i_1} \cdots X^{(n)}_{i_{n-1} j_n}\left(U^*_{j_n} \otimes P_{j_n}\right) Y_{j_n 1}\right) = 0$$

Similarly, we can get that

$$\tau\left(\left(U_2 \otimes P_2\right) X^{(1)}_{2j_1}\left(U^*_{j_1} \otimes P_{j_1}\right) Y_{j_1 i_1} \cdots X^{(n)}_{i_{n-1} j_n}\left(U^*_{j_n} \otimes P_{j_n}\right) Y_{j_n 2}\right) = 0$$

for $i_{1,} \cdots, i_{n-1}, j_{1,} \cdots, j_n \in \{1,2\}$. It follows that $\tau\left(A_{11}\right) = \tau\left(A_{22}\right) = 0$. Note that

$$\widetilde{E}\left(UX_1U^*Y_1UX_2U^*Y_2 \cdots UX_nU^*Y_n\right) = \sum_{i=1}^{2} \frac{1}{\tau\left(I \otimes P_i\right)}\left(\tau\left(A_{ii}\right) I \otimes P_i\right),$$

therefore

$$\widetilde{E}\left(UX_1U^*Y_1UX_2U^*Y_2 \cdots UX_nU^*Y_n\right) = 0$$

in $C^*_r\left(F_2\right) *_{red}\left(\mathcal{M}_{k_1}\left(\mathbb{C}\right) *_{red} \mathcal{M}_{k_2}\left(\mathbb{C}\right)\right)$.

Then, from the above argument, we conclude that $\left(\mathcal{M}_k\left(\mathbb{C}\right), E\right) \underset{\mathcal{D}}{*} \left(\mathcal{M}_k\left(\mathbb{C}\right), E\right)$ is an MF algebra.. ■

Although Proposition 3.1 is stated for a two-dimensional abelian C*-subalgebra, similar result holds for each finite-dimensional abelian C*-subalgebra by using a similar argument. We state this as a Theorem

Theorem 3.2 *Let $\mathcal{D} \subseteq \mathcal{M}_k\left(\mathbb{C}\right)$ be a unital inclusion of C*-algebras where $\mathcal{D}$ is a finite-dimensional abelian C*-algebra. Then, for the trace preserving conditional expectation $E: \mathcal{M}_k\left(\mathbb{C}\right) \to \mathcal{D}$, the reduced amalgamated free product $\left(\mathcal{M}_k\left(\mathbb{C}\right), E\right) \underset{\mathcal{D}}{*} \left(\mathcal{M}_k\left(\mathbb{C}\right), E\right)$ of $\left(\mathcal{M}_k\left(\mathbb{C}\right), E\right)$ and $\left(\mathcal{M}_k\left(\mathbb{C}\right), E\right)$ with respect to E is an MF algebra.*

Sketch of the proof We might assume that there are mutually orthogonal projections $P_1, \cdots, P_n$ in $\mathcal{M}_k\left(\mathbb{C}\right)$ such that

$$\mathcal{D} \cong \mathbb{C}P_1 \oplus \mathbb{C}P_2 \oplus \cdots \oplus \mathbb{C}P_n.$$

Let k_i =rank(P_i) for $1 \leq i \leq n$. Consider the unital C*-algebra:

$$\left(C^*_r\left(F_n\right) *_{red} \mathcal{M}_{k_1}\left(\mathbb{C}\right) *_{red} \cdots *_{red} \mathcal{M}_{k_n}\left(\mathbb{C}\right)\right) \otimes \mathcal{M}_k\left(\mathbb{C}\right).$$

Let $U_1, \cdots, U_n$ be the canonical generators of $C^*_r\left(F_n\right)$ and

$$U = U_1 \otimes P_1 + \cdots + U_n \otimes P_n$$

be a unitary in

$$\begin{aligned} & C_r^*(F_n) \otimes \mathcal{M}_k(\mathbb{C}) \\ \subseteq \quad & (C_r^*(F_n) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \cdots *_{red} \mathcal{M}_{k_n}(\mathbb{C})) \otimes \mathcal{M}_k(\mathbb{C}) . \end{aligned}$$

Similar argument in Lemma 3.5, shows that there is a unital C*-subalgebra $\mathcal{M}$ of $(\mathcal{M}_{k_1}(\mathbb{C}) *_{red} \cdots *_{red} \mathcal{M}_{k_n}(\mathbb{C})) \otimes \mathcal{M}_k(\mathbb{C})$ such that $\mathcal{M} \cong \mathcal{M}_k(\mathbb{C})$ and

$$\mathcal{M}_{k_1}(\mathbb{C}) \otimes P_1 + \cdots \oplus \mathcal{M}_{k_n}(\mathbb{C}) \otimes P_n \subseteq \mathcal{M}$$

as well as

$$\mathbb{C}(I \otimes P_1) \oplus \cdots \oplus \mathbb{C}(I \otimes P_n) \subseteq U\mathcal{M}U^*.$$

Let $C^*(\mathcal{M}, U\mathcal{M}U^*)$ be a unital C*-subalgebra generated by $\mathcal{M}$ and $U\mathcal{M}U^*$ in $(C_r^*(F_n) *_{red} \mathcal{M}_{k_1}(\mathbb{C}) *_{red} \cdots *_{red} \mathcal{M}_{k_n}(\mathbb{C})) \otimes \mathcal{M}_k(\mathbb{C})$. Then $C^*(\mathcal{M}, U\mathcal{M}U^*)$ is an MF algebra. Similar argument as Proposition 3.1 shows that

$$C^*(\mathcal{M}, U\mathcal{M}U^*) \cong (\mathcal{M}_k(\mathbb{C}), E) \underset{\mathcal{D}}{*} (\mathcal{M}_k(\mathbb{C}), E) .$$

This induces that $(\mathcal{M}_k(\mathbb{C}), E) \underset{\mathcal{D}}{*} (\mathcal{M}_k(\mathbb{C}), E)$ is also an MF algebra, which completes our proof. ■

3.3 $\mathcal{D}$ is a finite-dimensional C*-algebra

Now, we are ready to consider the case when $\mathcal{D}$ is a finite-dimensional C*-algebra.

Theorem 3.3 *Let $\mathcal{D} \subseteq \mathcal{M}_k(\mathbb{C})$ be a unital inclusion of C*-algebras where $\mathcal{D}$ is a finite-dimensional C^*-algebra and $E : \mathcal{M}_k(\mathbb{C}) \to \mathcal{D}$ be the trace preserving conditional expectation from $\mathcal{M}_k(\mathbb{C})$ onto $\mathcal{D}$. Then $(\mathcal{M}_k(\mathbb{C}), E) *_{\mathcal{D}} (\mathcal{M}_k(\mathbb{C}), E)$ is MF.*

Proof. Suppose $\mathcal{D} = \mathcal{M}_{k_1}(\mathbb{C}) \oplus \cdots \oplus \mathcal{M}_{k_t}(\mathbb{C})$ with $k_1 + \cdots + k_t = k$ and $P_1, \cdots, P_t$ are central projections of $\mathcal{D}$ with $\sum_{i=1}^t P_i = I \in \mathcal{M}_k(\mathbb{C})$. Let p_i be a minimal projection in $P_i\mathcal{D}$ for each $i \in \{1, \cdots, t\}$. Then $p_i \mathcal{D} p_i \cong \mathbb{C}p_i$. Assume $p = \sum_{i=1}^t p_i$, it follows that $p\mathcal{D}p$ is a finite-dimensional abelian C*-algebra. Note that there are partial isometries $v_1, \cdots, v_n \in \mathcal{D}$ with $v_i^* v_i \leq p$ and $\sum_{i=1}^n v_i v_i^* \leq 1 - p$ where $n = \max\{k_1, \cdots, k_t\} - 1$. So, by Lemma 3.1 and Lemma 3.2, it is sufficient to show that

$$\left(p\mathcal{M}_k(\mathbb{C})p, E|_{p\mathcal{M}_k(\mathbb{C})p}\right) *_{p\mathcal{D}p} \left(p\mathcal{M}_k(\mathbb{C})p, E|_{p\mathcal{M}_k(\mathbb{C})p}\right)$$

is an MF algebra. By Theorem 3.2, it is easy to see that

$$\left(p\mathcal{M}_k(\mathbb{C})p, E|_{p\mathcal{M}_k(\mathbb{C})p}\right) *_{p\mathcal{D}p} \left(p\mathcal{M}_k(\mathbb{C})p, E|_{p\mathcal{M}_k(\mathbb{C})p}\right)$$

is MF, so is $(\mathcal{M}_k(\mathbb{C}), E) *_{\mathcal{D}} (\mathcal{M}_k(\mathbb{C}), E)$. ■

It is not hard to extend the previous theorem to a reduced amalgamated free product of two finite-dimensional C*-algebras.

Theorem 3.4 *Let $\mathcal{M}_k(\mathbb{C}) \supseteq \mathcal{A} \supseteq \mathcal{D} \subseteq \mathcal{B} \subseteq \mathcal{M}_k(\mathbb{C})$ be unital C*-inclusions of finite-dimensional C*-algebras and $E : \mathcal{M}_k(\mathbb{C}) \rightarrow \mathcal{D}$ be the trace preserving conditional expectation. Then $(\mathcal{A}, E|_{\mathcal{A}}) *_{\mathcal{D}} (\mathcal{B}, E|_{\mathcal{B}})$ is MF.*

Proof. From the definition of reduced amalgamated free product of C*-algebras, we have the following C*-inclusion

$$(\mathcal{A}, E|_{\mathcal{A}}) *_{\mathcal{D}} (\mathcal{B}, E|_{\mathcal{B}}) \subseteq (\mathcal{M}_k(\mathbb{C}), E) *_{\mathcal{D}} (\mathcal{M}_k(\mathbb{C}), E).$$

So $(\mathcal{A}, E|_{\mathcal{A}}) *_{\mathcal{D}} (\mathcal{B}, E|_{\mathcal{B}})$ is an MF algebra. ■

Now we are ready to prove our main result in the paper.

Let $\mathbb{C}\langle \mathbf{X}_1, \ldots, \mathbf{X}_n\rangle$ be the set of all noncommutative polynomials in the indeterminants $\mathbf{X}_1, \ldots, \mathbf{X}_n$. Then we let

$$\mathbb{C}\langle \mathbf{X}_1, \mathbf{X}_2, \cdots\rangle = \cup_{m=1}^{\infty} \mathbb{C}\langle \mathbf{X}_1, \mathbf{X}_2, \cdots \mathbf{X}_m\rangle.$$

The following theorem give a necessary and sufficient condition such that a reduced free product of two UHF algebras with amalgamation over a finite-dimensional C*-algebra is MF.

Theorem 3.5 *Suppose that $\mathcal{A}_1$ and $\mathcal{A}_2$ are two unital UHF-algebras with the faithful tracial states $\tau_{\mathcal{A}_1}$ and $\tau_{\mathcal{A}_2}$ respectively. Let $\mathcal{A}_1 \supseteq \mathcal{D} \subseteq \mathcal{A}_2$ be unital inclusions of C*-algebras where $\mathcal{D}$ is a finite-dimensional C*-algebra. Assume that $E_{\mathcal{A}_1} : \mathcal{A}_1 \rightarrow \mathcal{D}$ and $E_{\mathcal{A}_2} : \mathcal{A}_2 \rightarrow \mathcal{D}$ are the trace preserving conditional expectations onto $\mathcal{D}$. Then the reduced amalgamated free product $(\mathcal{A}_1, E_{\mathcal{A}_1}) *_{\mathcal{D}} (\mathcal{A}_2, E_{\mathcal{A}_2})$ is an MF algebra if and only if $\tau_{\mathcal{A}_1}(z) = \tau_{\mathcal{A}_2}(z)$ for all $z \in \mathcal{D}$.*

Proof. From the fact that every MF algebra has a tracial state and $\mathcal{A}_1$, $\mathcal{A}_2$ have unique tracial states, one direction of the proof is obvious. Suppose $\mathcal{A}_1 = \overline{\cup_{k=1}^{\infty} \mathcal{M}_{n_k}(\mathbb{C})}$ and $\mathcal{A}_2 = \overline{\cup_{l=1}^{\infty} \mathcal{M}_{m_l}(\mathbb{C})}$ where $\{\mathcal{M}_{n_k}(\mathbb{C})\}_{k=1}^{\infty}$ and $\{\mathcal{M}_{m_l}(\mathbb{C})\}_{l=1}^{\infty}$ are increasing sequences of full matrix algebras. Since $\mathcal{D}$ is a finite-dimensional C*-algebra, without loss of generality, we may assume that there is an integer t_0 such that, for $\forall t \geq t_0$, we have $\mathcal{D} \subseteq \mathcal{M}_{n_t}(\mathbb{C})$ and $\mathcal{D} \subseteq \mathcal{M}_{m_t}(\mathbb{C})$. From the equation $\tau_{\mathcal{A}}(z) = \tau_{\mathcal{B}}(z)$ for all $z \in \mathcal{D}$, there is an integer s such that $\mathcal{M}_{n_t}(\mathbb{C})$ and $\mathcal{M}_{m_t}(\mathbb{C})$ can be both unital embedded into $\mathcal{M}_s(\mathbb{C})$ satisfying

$$\mathcal{M}_s(\mathbb{C}) \supseteq \mathcal{M}_{n_t}(\mathbb{C}) \supseteq \mathcal{D} \subseteq \mathcal{M}_{m_t}(\mathbb{C}) \subseteq \mathcal{M}_s(\mathbb{C}).$$

Therefore, by Theorem 3.4, we have that

$$\left(\mathcal{M}_{n_t}(\mathbb{C}), E_{\mathcal{A}_1}|_{\mathcal{M}_{n_t}(\mathbb{C})}\right) *_{\mathcal{D}} \left(\mathcal{M}_{m_t}(\mathbb{C}), E_{\mathcal{A}_2}|_{\mathcal{M}_{m_t}(\mathbb{C})}\right)$$

is an MF algebra.

Suppose $\{x_1, x_2, \cdots, z_1, \cdots z_h\}$ is a family of self-adjoint generators of $\mathcal{A}_1$ with $\|x_i\| = \|z_j\| = 1$, $\{y_1, y_2, \cdots z_1, \cdots, z_h\}$ is a family of self-adjoint generators of $\mathcal{A}_2$ with $\|y_i\| = \|z_j\| = 1$ for all $i \in \mathbb{N}$ and $j \in \{1, \cdots, h\}$ where $\{z_1, \cdots, z_h\}$ is a family of self-adjoint generators of $\mathcal{D}$. For any $\varepsilon > 0$ and any $r \in \mathbb{N}$ with a family of noncommutative polynomials $P_1, \cdots, P_r$ in

$$\mathbb{C}\langle X_1, \cdots X_r, Y_1, \cdots Y_r, Z_1, \cdots, Z_h\rangle,$$

there is a integer $t \geq t_0$ with

$$T_1, \cdots, T_r \in \mathcal{M}_{m_t}^{s.a.}(\mathbb{C}) \subseteq \left(\mathcal{M}_{n_t}(\mathbb{C}), E_{\mathcal{A}_1}|_{\mathcal{M}_{n_t}(\mathbb{C})}\right) *_{\mathcal{D}} \left(\mathcal{M}_{m_t}(\mathbb{C}), E_{\mathcal{A}_2}|_{\mathcal{M}_{m_t}(\mathbb{C})}\right),$$

$$S_1, \cdots, S_r \in \mathcal{M}_{n_t}^{s.a.}(\mathbb{C}) \subseteq \left(\mathcal{M}_{n_t}(\mathbb{C}), E_{\mathcal{A}_1}|_{\mathcal{M}_{n_t}(\mathbb{C})}\right) *_{\mathcal{D}} \left(\mathcal{M}_{m_t}(\mathbb{C}), E_{\mathcal{A}_2}|_{\mathcal{M}_{m_t}(\mathbb{C})}\right)$$

and

$$K_1, \cdots, K_h \in \mathcal{D} \subseteq \left(\mathcal{M}_{n_t}(\mathbb{C}), E_{\mathcal{A}_1}|_{\mathcal{M}_{n_t}(\mathbb{C})}\right) *_{\mathcal{D}} \left(\mathcal{M}_{m_t}(\mathbb{C}), E_{\mathcal{A}_2}|_{\mathcal{M}_{m_t}(\mathbb{C})}\right)$$

such that, for $1 \leq j \leq r$,

$$\begin{aligned} & |\|P_j(x_1, \cdots, x_r, y_1, \cdots, y_r, z_1, \cdots, z_h)\| - \|P_j(T_1, \cdots, T_r, S_1, \cdots, S_r, K_1, \cdots, K_h)\|| \\ \leq \ & \|P_j(x_1, \cdots, x_r, y_1, \cdots, y_r, z_1, \cdots, z_h) - P_j(T_1, \cdots, T_r, S_1, \cdots, S_r, K_1, \cdots, K_h)\| \\ < \ & \varepsilon/2 \end{aligned}$$

where

$$P_j(T_1, \cdots, T_r, S_1, \cdots, S_r, K_1, \cdots, K_h) \in \left(\mathcal{M}_{n_t}(\mathbb{C}), E_{\mathcal{A}_1}|_{\mathcal{M}_{n_t}(\mathbb{C})}\right) *_{\mathcal{D}} \left(\mathcal{M}_{m_t}(\mathbb{C}), E_{\mathcal{A}_2}|_{\mathcal{M}_{m_t}(\mathbb{C})}\right).$$

Since $\left(\mathcal{M}_{n_t}(\mathbb{C}), E_{\mathcal{A}_1}|_{\mathcal{M}_{n_t}(\mathbb{C})}\right) *_{\mathcal{D}} \left(\mathcal{M}_{m_t}(\mathbb{C}), E_{\mathcal{A}_2}|_{\mathcal{M}_{m_t}(\mathbb{C})}\right)$ is MF, there is a positive integer k and self-adjoint matrices $A_1, \cdots, A_r, B_1, \cdots, B_r, C_1, \cdots, C_t \in \mathcal{M}_k^{s.a.}(\mathbb{C})$ such that, for all $1 \leq j \leq r$,

$$|\|P_j(A_1, \cdots, A_r, B_1, \cdots, B_r, C_1, \cdots, C_h)\| - \|P_j(T_1, \cdots, T_r, S_1, \cdots, S_r, K_1, \cdots, K_h)\|| \leq \varepsilon/2.$$

This implies that

$$|\|P_j(x_1, \cdots, x_r, y_1, \cdots, y_r, z_1, \cdots, z_h)\| - \|P_j(A_1, \cdots, A_r, B_1, \cdots, B_r, C_1, \cdots, C_h)\|| < \varepsilon.$$

Then, by Lemma 2.4.1 in [16], $(\mathcal{A}_1, E_{\mathcal{A}_1}) *_{\mathcal{D}} (\mathcal{A}_2, E_{\mathcal{A}_2})$ is MF. ■

We end this subsection with a result on the reduced C*-algebra of an amalgamated free product group of two finite groups. First, we need the following lemma.

Lemma 3.6 *Suppose that $A \supseteq H \subseteq B$ are fnite groups satisfying $[A : H] \geq 3$ and $[B : H] \geq 2$. Let $A *_H B$ be the amalgamated free product of groups A and B over H. Then $A *_H B$ is not amenable.*

Proof. By Corollary 4.10 of [6], we know that

$$\delta_0(\mathbb{C}(A *_H B)) = \left(1 - |A|^{-1}\right) + \left(1 - |B|^{-1}\right) - \left(1 - |H|^{-1}\right) = 1 + \left(|H|^{-1} - |A|^{-1} - |B|^{-1}\right)$$

where δ_0 is the Voiculescu's (microstate) modifed free entropy dimension and $|A|$, $|B|$ and $|H|$ are the cardinalities of groups A, B and H respectively. Note that

$$|H|^{-1} - |A|^{-1} - |B|^{-1} = |H|^{-1}\left(1 - \frac{1}{[A : H]} - \frac{1}{[B : H]}\right) \geq |H|^{-1}\left(1 - \frac{1}{3} - \frac{1}{2}\right) > 0.$$

Hence $\delta_0(\mathbb{C}[A *_H B]) > 1$. By Corollary 6.8 in [18], we know that $A *_H B$ is not an amenable group. ■

Applying preceding Theorem and Lemma, we obtain the following corollary.

Corollary 3.1 *Suppose that $A \supseteq H \subseteq B$ are fnite groups and $A *_H B$ is the amalgamated free product of groups A and B over H. Then $C_r^*(A *_H B)$ is an MF algebra. If futher $[A : H] \geq 3$ and $[B : H] \geq 2$, then Ext $(C_r^*(A *_H B))$, the Billmore-Douglus-Fillmore extension semigroup, is not a group.*

Proof. Note that

$$C_r^*(A *_H B) \cong (C_r^*(A), E_A) *_{C_r^*(H)} (C_r^*(B), E_B)$$

where E_A and E_B are canonical trace preserving conditional expectations from $C_r^*(A)$ onto $C_r^*(H)$, and from $C_r^*(B)$ onto $C_r^*(H)$ respectively (see [8]). Now it follows from Theorem 3.5 that $C_r^*(A *_H B)$ is an MF algebra. Moreover, if $[A : H] \geq 3$ and $[B : H] \geq 2$, by Lemma 3.6, we get that $A *_H B$ is not amenable, whence $C_r^*(A *_H B)$ is not a quasidiagonal C* -algebra. Recall a well-known fact that if a unital C -algebra $\mathcal{E}$ is an MF algebra but not a quasidiagonal C* -algebra, then Ext($\mathcal{E}$) is not a group. Therefore, Ext($C_r^*(A *_H B)$), the Billmore-Douglus-Fillmore extension semigroup, is not a group.

■

References

[1] D. Avitzour, Free Products of C*-algebras, Trans. Amer. Math. Soc. 271, 1982, 423-435.

[2] William. Arveson, An invitation to C*-algebras, Graduate Texts in Mathematics, No. 39, Springer-Verlag, New York-Heidelberg, 1976.

[3] B. Blackadar, Operator algebras. Theory of $Cˆ*$-algebras and von Neumann algebras. Encyclopaedia of Mathematical Sciences, 122. Operator Algebras and Non-commutative Geometry, III. Springer-Verlag, Berlin, 2006.

[4] B. Blackadar and E. Kirchberg, Generalized inductive limits of finite dimensional C*-algebras, Math. Ann. 307 (1997), 343-380.

[5] N. P. Brown and K. J. Dykema, Popa algebras in free group factors, J. reine angew. Math 573 (2004), 157-180.

[6] N. P. Brown, K. J. Dykema and K. Jung, Free entropy dimension in amalgamated free products, Proc. London Math. Soc. (3) 97 (2008) 339–367

[7] E. F. Blanchard and K. J. Dykema, Embeddings of reduced free products of operator algebras, Pacific Journal of Mathematics Volume 199(2001), No. 1, 1-19.

[8] N. P. Brown and N. Ozawa, Algebras and Finite-Dimensional Approximations, American Mathematical Society (March 12, 2008).

[9] K. Davidson, C*-algebras by Example, Amer Mathematical Society (June 1996).

[10] K. J. Dykema, A description of amalgamated free products of finite von Neumann algebras over finite dimensional subalgebras, arXiv:0911.2052v3

[11] K. J. Dykema, Exactness of reduced amalgamated free product C*-algebras, Forum Math. 16 (2004), 161-180.

[12] K. J. Dykema and D. Shlyakhtenko, exactness of Cuntz-Pimsner C*-algebras, Proc. Edinb. Math. Soc. 44 (2001), 425-444.

[13] U. Haagerup, S. Thorbjϕrnsen, A new application of random matrices: $Ext(C^*_{red}(F_2))$ is not a group, Ann. of Math. (2) 162 (2005), no. 2, 711–775.

[14] D. Hadwin, Q. Li and J. Shen, "Topological free entropy dimensions in Nuclear C*-algebras and in Full Free Products of Unital C*-algebras".

[15] D. Hadwin; J. Shen, Some Examples of Blackadar and Kirchberg's MF Algebras, arXiv:0806.4712v4

[16] D. Hadwin; J. Shen, Reduced Free Products of Unital AH Algebras and Blackadar and Kirchberg's MF Algebras, arXiv:0812.0189v1

[17] N. A. Ivanov, On the structure of some reduced amalgamated free product C*-algebras, arXiv:0705.3919.

[18] K. Jung, The free entropy dimension of hyperfinite von Neumann algebras, Trans. Amer. Math. Soc. 355 (2003), 5053-5089.

[19] R. Kadison, J. Ringrose, Fundamentals of the Operator Algebras (Academic, Orlando, FL), (1983, 1986)Vols. 1 and 2.

[20] I. Kaplansky, The structure of certain operator algebras, Trans. Amer. Math. Soc. 70 (1951),219-255.

[21] E. C. Lance, Hilbert C*-modules: a toolkit for operator algebraists(London Mathematical Society Lecture Note Series), Cambridge University Press (March 31, 1995).

[22] D. Voiculescu, The topological version of free entropy, Lett. Math. Phys. 62 (2002), no. 1, 71–82.

[23] D. Voiculescu, Symmetries of some reduced free product C*-algebras, Operator Algebras and Their Connection with Topology and Ergodic Theory, Lecture Notes in Mathematics, Volume 1132, Spring-Verlag, 1985, 556-588.

Junhao Shen
Department of Mathematics and Statistics
Univeristy of New Hampshire
Durham, NH, 03824
Email: jog2@cisunix.unh.edu

Qihui Li
Department of Mathematics and Statistics
Univerisity of New Hampshire
Durham, NH, 03824
Email: qme2@cisunix.unh.edu

and
Department of Mathematics
East China University of Science and Technology
Shanghai, China Meilong Road 130, 200237